

Osteochondritis Dissecans (OCD) Repair

David E. Hartigan, M.D.

Patient Name	Date
--------------	------

Procedure: Right Left Knee Arthroscopy OCD repair

Associated Procedure: (check if applicable) OATS Bone graft harvest (tibia, ICBG) Osteotomy (DFO, HTO, or AMZ)

Evaluate and Treat – no open chain or isokinetic exercises

Provide patient with home exercise program

Phase I (Weeks 0-8)*:** *Period of protection. NWB with crutches until after hardware is removed (2nd operation). Progress ROM as tolerated.*

- **Non-weight bearing with crutches.** Hinged knee brace should be worn at all times except during PT and for hygiene. Brace should be locked in extension when not ambulating or performing PT. For patients without CPM at night, brace should be locked in extension while sleeping.
- **ROM:** Progress through passive, active and active-assisted ROM as tolerated
 - Goal: Full extension by 2 weeks, 130 degrees of flexion by 6 weeks
- Patellar mobilization as tolerated.
- **Strengthening:** quad sets, SLRs (with brace locked in extension), heel slides, etc. Ankle/hip strengthening as tolerated.

****In some situations, a CPM device will be ordered for home use: 6-8 hours per day (usually at night) x 6 weeks; start at 0-40 degrees (1 cycle/minute), progressing 5-10 degrees daily as tolerated (goal 120 degrees or greater by week 6). For those without CPM, the brace should be locked in extension during sleep.*

Other:

Modalities

Ultrasound

Heat before/after

Ice before/after exercise

Electrical Stimulation

Frequency: _____ x/week x _____ weeks

By signing this referral, I certify that I have examined this patient and physical therapy is medically necessary.

This patient ____ would ____ would not benefit from social services.

Physician Signature	Date
Printed Name	